

- Cigarette smoking is the single most preventable cause of death in Hawai'i, [1] as well as in the United States, and has been called the nation's "number one" public health problem. [2]
- More than 400,000 Americans [2] and 1200 people in Hawai'i [3] die each year from tobacco related causes.
- Smokers lose an average of 12 to 15 years of life. [4]
- Cigarette smoking causes more premature deaths each year,
 (before the age of 65), than all of the following causes of death together:
 AIDS, drug abuse, alcohol, and all unintentional injury (motor vehicle, drowning and falls) as well as all intentional injury (suicide and homicide). [5]



Data Source: Behavior Risk Factor Surveillance Survey: Hawai'i State Dept. of Health



Data Source: Office of Health Status Monitoring, Hawaii State Dept. of Health

REFERENCE:

- 1. Centers for Disease Control and Prevention. (2001). Investment in Tobacco Control: State Highlights 2001. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health
- 2. U.S. Department of Health and Human Services. (2000). Reducing Tobacco Use: A Report of the Surgeon General. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- 3. Department of Health, Vital Records.
- 4. The Center for Social Gerontology. (1999). Fact Sheet on Tobacco & Older Persons. Available: www.tcsq.org/tobacco/facts.htm [2001, 6/1/01].
- 5. HIV/AIDS Surveillance Report; National Safety Council Accident Sheets; Monthly Vital Statistics Report; SAMMEC; Alcohol-Related Death Index.